

IN THE CLAIMS

I claim

- [C1]** A method of stimulating an immune response in a virally infected individual, the method comprising:
- providing an imidazoquinolinamine formulation;
 - disposing an amount of the imidazoquinolinamine formulation into a first nare of a virally infected individual; and
 - covering at least a portion of the internal surface of the individual's first nare with a portion of the amount of the imidazoquinolinamine in said nare.
- [C2]** The method of claim 1, wherein the imidazoquinolinamine formulation's active ingredient is selected from the group consisting of imiquimod and resiquimod.
- [C3]** The method of claim 1, wherein the imidazoquinolinamine formulation is disposed using a device selected from the group consisting of a swab, syringe, spray nozzle, and a drip applicator.
- [C4]** The method of claim 1, wherein the imidazoquinolinamine formulation comprises imiquimod as an active ingredient in the amount of about 12 mg.
- [C5]** The method of claim 1, wherein the imidazoquinolinamine formulation comprises resiquimod as an active ingredient in the amount of about 12 mg.
- [C6]** The method of claim 1, wherein the imidazoquinolinamine formulation is selected from a group consisting of a cream, gel, liquid, paste, aerosol, and an emulsion.
- [C7]** The method of claim 1, wherein about 15 minutes prior to disposing the imidazoquinolinamine formulation, the first nare is treated with a decongestant.
- [C8]** The method of claim 7, wherein the decongestant is Neosynepherine.
- [C9]** The method of claim 1, wherein the individual is infected with viral rhinitis.
- [C10]** The method of claim 1 further comprising providing the imidazoquinolinamine formulation every twelve hours until relief from cold symptoms is obtained.
- [C11]** The method of claim 1, wherein the imidazoquinolinamine formulation is applied within twenty four hours after an appearance of first cold symptoms.